

Search for Interstellar Water in the Translucent Molecular Cloud toward HD 154368

Marco Spaans¹, David Neufeld

Department of Physics & Astronomy, The Johns Hopkins University, Baltimore, MD 21218

Stephen Lepp

University of Nevada, 4505 S. Maryland Parkway, Las Vegas, NV 89154-4002

Gary J. Melnick, and John Stauffer

Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

ABSTRACT

We report an upper limit of $9 \times 10^{12} \text{ cm}^{-2}$ on the column density of water in the translucent cloud along the line-of-sight toward HD 154368. This result is based upon a search for the C-X band of water near $1240 \text{ \AA}$ carried out using the Goddard High Resolution Spectrograph of the Hubble Space Telescope. Our observational limit on the water abundance together with detailed chemical models of translucent clouds and previous measurements of OH along the line-of-sight constrain the branching ratio in the dissociative recombination of H_3O^+ to form water. We find at the 3σ level that no more than 30% of dissociative recombinations of H_3O^+ can lead to H_2O .

The observed spectrum also yielded high-resolution observations of the Mg II doublet at $1239.9 \text{ \AA}$ and $1240.4 \text{ \AA}$, allowing the velocity structure of the dominant ionization state of magnesium to be studied along the line-of-sight. The Mg II spectrum is consistent with GHRS observations at lower spectral resolution that were obtained previously but allow an additional velocity component to be identified.

Subject headings: Ultraviolet: ISM: Lines and Bands – ISM: Molecules – Molecular Processes – Ultraviolet: ISM

1. Introduction

Translucent clouds have total extinctions A_V in the range of 2-5 magnitudes. The physical and chemical conditions that characterize translucent clouds are therefore intermediate between those in diffuse and in dense molecular clouds (Crutcher 1985; van Dishoeck & Black 1988). Photodissociation rates are significantly smaller in translucent clouds than in diffuse clouds, and the column densities of molecules like CO, OH and CS are correspondingly larger. Translucent clouds can be observed through absorption lines of CN, CH, CH^+ and C_2 as well as through millimeter emission lines of CO and CS. Although they are an abundant constituent of dense molecular clouds (Jacq et al. 1988; Zmuidzinas et al. 1995; Gensheimer et al. 1996; van Dishoeck & Helmich 1996; Cernicharo et al. 1997) and are expected to be the dominant coolant of warm dense regions within such clouds (Neufeld & Kaufman 1993; Neufeld, Lepp & Melnick 1995), water

¹Hubble Fellow, currently at the Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138

molecules have not been detected in diffuse or translucent molecular clouds. In this paper, we report the results of a search for water in the translucent cloud along the line of sight to HD 154368.

HD 154368 is an O9.5 Iab star at an estimated distance of 800 pc from the Sun (Snow et al. 1996) that is situated near the Sco OB 1 association (Blades & Bennewith 1973). The line-of-sight toward HD 154368 lies about 14° from the core of the ρ Oph molecular cloud, which is located at a distance of 125 ± 25 pc (de Geus, de Zeeuw, & Lub 1989) and has a heliocentric radial velocity of -6.6 km s^{-1} . The main H I (e.g. Riegel & Crutcher 1972) and Na I (e.g. Crawford, Barlow, & Blades 1989) absorption features are observed at a radial velocity that is close to that of the ρ Oph molecular cloud, a result that suggests that the main component of gas toward HD 154368 is probably not close to the star and is more likely to be the outer envelope of a dense molecular cloud only about 125 pc from the Sun. The color excess along the line of sight is $E(B-V)=0.82$ and most of the gas toward HD 154368 resides in two clouds centered near -3.26 (main component) and -20.95 km s^{-1} (heliocentric).

This line-of-sight has been observed extensively from the ground by means of narrow H I 21 cm absorption features (e.g. Riegel & Crutcher 1972); optical absorption lines of Na I D (e.g. Crawford et al. 1989), CN B-X (0,0) and A-X (0,0), CH, CH^+ ; and near-infrared absorption lines of C_2 in the A-X Phillips system at $8750 \text{ \AA}$ (van Dishoeck & de Zeeuw 1984). The CH observations can be used to infer the total column density of H_2 along the line-of-sight (Gredel, van Dishoeck, & Black 1993), a quantity that is needed to determine the atomic depletions. The red system of CN has been used by Gredel, van Dishoeck, & Black (1991) to derive an electron density of $0.05\text{--}0.15 \text{ cm}^{-3}$ for the molecular component. Molecular emission lines are also detectable toward HD 154368. Data on $^{12}\text{CO } J = 1 - 0$, $J = 2 - 1$, and $J = 3 - 2$ and $^{13}\text{CO } J = 1 - 0$ have been presented by van Dishoeck et al. (1991), and have been used to constrain the column density of CO and the density of H_2 in the molecular component. These authors confirmed the relatively low density $n_{\text{H}} \approx 350 \text{ cm}^{-3}$ derived independently from the C_2 absorption data. The $^{12}\text{CO } J = 1 - 0$ distribution over a region of $30' \times 30'$ around the star is featureless, a result that suggests once more that the cloud and the star are not located close to one another.

The velocity structure of the gas toward HD 154368 is well known through the Na I data obtained using the Ultra High Resolution Facility at the Anglo Australian Telescope (Snow et al. 1996) and high resolution Ca II observations (Crawford 1992). The Na I data indicates seven velocity components at -27.7 , -20.95 , -18.2 , -14.75 , -10.5 , -3.26 , and $+5.62 \text{ km s}^{-1}$ where 96% of the neutral sodium resides in the -3.26 km s^{-1} feature. The Ca II results show five velocity components at -27.6 , -20.9 , -14.4 , -4.3 , and $+6.8 \text{ km s}^{-1}$. These five coincide roughly with the Na I components, but their relative column densities are different. Approximately 50% of the ionized calcium resides in the -4.3 km s^{-1} feature with an even distribution over the remaining velocity components.

A detailed discussion of all these observations and the implications they have for the physical conditions of the ambient medium can be found in Snow et al. (1996). In this work, their results will be adopted and the main focus will be on the resulting OH and H_2O chemistry.

2. Observational Results

Observations of HD 154368 in the $1234.49\text{--}1241.35 \text{ \AA}$ spectral region were obtained at a spectral resolution of $\sim 3 \text{ km s}^{-1}$ using Echelle-A of the Goddard High Resolution Spectrograph (GHRS) on HST. The data, obtained with the use of standard substepping, were reduced within the IRAF environment

following procedures developed by the GHRS Science Team (Cardelli, Ebbets, & Savage 1990; 1993). The equivalent widths and wavelengths of the features in the calibrated spectra were obtained with various IRAF routines and in particular the multi-component “specfit” package developed by Kriss (1994) that allows for deblending.

Figures 1a and 1b show the region around the C 1B_1 -X 1A_1 band of water near 1240 Å. We find no evidence in the spectrum for features at the expected location of either the ortho lines $1_{01} - 2_{11}$ (1239.382 Å) and $1_{01} - 1_{11}$ (1240.153 Å), or the para line $0_{00} - 1_{10}$ (1239.728 Å). To determine a reliable upper limit in the presence of noise, an artificial water feature was added until a 3σ identification could be made with the specfit program. The assumed line width for the artificial feature was $b = 2.9 \text{ km s}^{-1}$, the measured b -value for the dominant MgII component (see Table 1 below). With the water oscillator strengths of Smith & Parkinson (1978) and Smith et al. (1981), and the assumptions that one-quarter and three-quarters of the water molecules are respectively in the 0_{00} and 1_{01} states, we find a 3σ upper limit of $9 \times 10^{12} \text{ cm}^{-3}$ for the column density of water toward HD 154368. This upper limit was obtained by co-adding the separate lines in the spectrum, and was not found to change when the more recent line rest wavelengths of Watson (1997, private communication) and the oscillator strengths of Watson (1997, private communication) and Yoshino (1997, private communication) were used. The 3σ upper limits on the equivalent widths of the 1239.382 Å, 1239.728 Å and 1240.153 are 0.85 mÅ, 0.89 mÅ and 0.84 mÅ, respectively. For comparison, the limit on the H₂O column density in the more diffuse ($A_V \approx 1 \text{ mag}$) ζ Oph interstellar cloud is $3 \times 10^{12} \text{ cm}^{-2}$ (Snow 1976).

Figure 2 shows the Mg II doublet features around 1239.9 Å and 1240.4 Å. Table 1 presents the results of the fitting procedure for Mg II and lists the heliocentric velocities of the various components, their b values, equivalent widths and column densities. The fit is based upon the line rest wavelengths of Morton (1991) and the oscillator strengths of Fitzpatrick (1997). Our results for the equivalent widths are in good general agreement with those obtained previously from lower resolution spectra of Mg II (Snow et al. 1996), although our derived column densities are a factor of 2.4 smaller due to the use of the Fitzpatrick (1997) oscillator strengths, rather than those tabulated by Morton (1991). A systematic velocity shift of $\sim 2 \text{ km s}^{-1}$ is present. This shift is not inconsistent with the GHRS wavelength calibration uncertainty and is most likely caused by geo-magnetic effects. We also find an additional velocity feature at -36.2 km s^{-1} , apparent in both components of the doublet, with a column density $\sim 1 \times 10^{14} \text{ cm}^{-3}$, which has not been detected previously.

Figure 3 shows a clear detection of the Ge II transition at 1237.059 Å. The heliocentric velocity of this component is -1.2 km s^{-1} , in excellent agreement with that of the main Mg II component. No other GeII features fall in the spectral region covered by our observations.

3. Model

To interpret our upper limit on the water column density toward HD 154368, we used the numerical code described by Spaans (1996) and by Spaans & van Dishoeck (1997) to construct chemical models of the foreground cloud. A detailed description of the numerical model is given in these references and the interested reader is referred there for details. The model input consists of the density (assumed constant) n_H , the total extinction through the cloud A_V , the gas phase elemental abundances, the strength of the illuminating radiation field I_{UV} in units of the average interstellar radiation field of the Galaxy, the

shape of the ultraviolet extinction curve, the cosmic ray ionization rate ζ , and the branching ratios for the dissociative recombination of H_3O^+ . The cloud is modeled as a two-sided plane parallel slab. Dust attenuation and self-shielding of H_2 and CO are included explicitly through a Monte Carlo radiative transfer method. The extinction curve of Snow et al. (1996) is adopted for the specific line of sight toward HD 154368, and is extended to wavelengths of 1000 Å using the method of Cardelli, Clayton, & Mathis (1989) with a value of the ratio of total to selective extinction of $R_V = 3.1$.

The model produces as output the thermal and chemical structure of the cloud and the column densities of all major chemical species. Figures 4a and 4b present the results for a model with $n_{\text{H}} = 325 \text{ cm}^{-3}$, $I_{\text{UV}} = 3$, $\zeta = 5 \times 10^{-17} \text{ s}^{-1}$, a H_3^+ recombination rate of $8 \times 10^{-7} T^{-0.7} \text{ cm}^3 \text{ s}^{-1}$ (Smith & Spañel 1993) and the depletions as given in Snow et al. (1996), which reproduces the observations to within 50% for all observed species except CH^+ . The CH^+ problem is well known in interstellar studies and no attempt is made here to alleviate it (e.g. Dalgarno 1976; Black 1988). The C_2 observations of van Dishoeck & de Zeeuw (1984) yield a kinetic gas temperature of 25 K in the molecular region of the cloud ($A_V > 1 \text{ mag}$). The temperature dependence depicted in Figure 4b is consistent with this value. The dust temperature is assumed to be constant and does not strongly influence the chemistry.

The predicted abundances of water and OH depend critically upon two poorly-known parameters to which the abundances of other observed species are insensitive: the cosmic ray ionization rate, ζ , and the branching ratio for dissociative recombination of H_3O^+ . We define $f(\text{H}_2\text{O})$, $f(\text{OH})$, and $f(\text{O})$ as the fractions of dissociative recombinations of H_3O^+ that lead to H_2O , OH and O.

With the cloud geometry, density and radiation field fixed by observations of other species, one can construct a grid of models with different values of ζ and $f(\text{H}_2\text{O})$. The OH and H_2O column densities predicted in these simulations are shown in Figure 5 (solid lines). For these models we have assumed that $f(\text{O})=0$ (or equivalently, $f(\text{H}_2\text{O}) + f(\text{OH}) = 1$), as has been assumed in most theoretical studies and is motivated by the recent experiment of Vejby-Christensen et al. (1997). The cross indicates the 3σ limit on the water column density reported here and the observed OH column density of $N(\text{OH})=1.4 \times 10^{14} \text{ cm}^{-2}$ (Black 1997, private communication). From these results it follows at the 3σ confidence level that $f(\text{H}_2\text{O})$ can be no larger than 0.30 and ζ no larger than $1 \times 10^{-16} \text{ s}^{-1}$. This rules out the large $f(\text{H}_2\text{O})$ branching ratio ~ 1 predicted by Bates (1986; but note the revised estimates of Bates 1991), provided that our adopted water chemistry, and in particular the value of the H_3^+ recombination rate, is secure.

Two recent experiments have yielded contradictory results for the H_3O^+ branching ratios. The experiment of Vejby-Christensen et al. (1997) implied that $f(\text{OH}) = 0.66$ and $f(\text{H}_2\text{O}) = 0.33$, values that are formally ruled out by our limit on the water column density, at least given the model assumptions discussed above. The results of Williams et al. (1996), by contrast, suggested that the dissociative recombination of H_3O^+ has a significant channel to O ($f(\text{O}) = 0.30$) and that the channel to H_2O is much less important ($f(\text{H}_2\text{O}) = 0.05$). To investigate the influence of the channel to O, we have run additional models in which $f(\text{H}_2\text{O}) + f(\text{OH}) = 0.7$ rather than unity. The dashed curve in Figure 5 denotes the results for these computations with $f(\text{H}_2\text{O}) = 0.05$. For $f(\text{O}) = 0.3$, our observed 3σ upper limit on the water column density is evidently consistent with the results of Williams et al. (and indeed with any value of $f(\text{H}_2\text{O}) \leq 0.15$). Our results leave open the possibility that both laboratory results are consistent, but that some water measured by Vejby-Christensen et al. (1997) is in a pre-dissociated state (Adams 1997, private communication).

4. Summary

We have reported a 3σ upper limit of $9 \times 10^{12} \text{ cm}^{-2}$ on the column density of water toward HD 154368. We have constructed detailed chemical models which incorporate many existing constraints on the physical conditions of the ambient medium. Together with the known column density of OH along the line of sight, we constrain the H_2O branching ratio for dissociative recombination of H_3O^+ to be smaller than 30%. Our results are in agreement with the laboratory studies of Williams et al. which find an oxygen channel in the recombination of H_3O^+ and a small branching ratio for the production of water, but are mildly inconsistent with the laboratory results of Vejby-Christensen et al.

The observed spectrum of HD 154368 also yielded high-resolution observations of the Mg II doublet at 1239.9 Å and 1240.4 Å, allowing the velocity structure of the dominant ionization state of magnesium to be studied along the line-of-sight. The Mg II spectrum is consistent with GHRS observations at lower spectral resolution that were obtained previously but allow an additional velocity component to be identified. Absorption by the Ge II line at 1237.059 Å was also detected, the first detection of interstellar germanium along the line-of-sight to HD 154368.

We acknowledge with gratitude the support of NASA grant NAGW-3147 from the Long Term Space Astrophysics Research Program and of grant GO-06739.01-95A from the HST Cycle 6 Program. In the final stages of this project MS was partially supported by NASA through grant HF-01101.01-97A, awarded by the Space Telescope Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA under contract NAS 5-26555. Most of the simulations presented in this work were performed on the Cray YMP operated by the Netherlands Council for Supercomputer Facilities in Amsterdam.

REFERENCES

- Bates, D.R. 1991, *J. Phys. B*, 24, 3267
- Bates, D.R. 1986, *ApJ*, 308, L45
- Black, J.H. 1988, *Adv. Atom. Mol. Phys.*, 25, 477
- Blades, J.C., & Bennewith, P.D. 1973, *MNRAS*, 161, 213
- Cardelli, J.A., Ebbets, D.C., & Savage, B.D. 1990, *ApJ*, 365, 789
- Cardelli, J.A., Ebbets, D.C., & Savage, B.D. 1993, *ApJ*, 413, 401
- Cardelli, J.A., Clayton, G.C., & Mathis, J.S. 1989, *ApJ*, 345, 245
- Cernicharo, J., et al. 1997, *A&A*, 323, L25
- Crawford, I.A. 1992, *MNRAS*, 259, 47
- Crawford, I.A., Barlow, M.J., & Blades, J.C. 1989, *ApJ*, 336, 212
- Crutcher, R.M. 1985, *ApJ*, 254, 82
- Dalgarno, A. 1976, in *Atomic Processes and Applications*, eds. P.G. Burke and B.L. Moiseiwitsch (Amsterdam: North-Holland), p. 110
- de Geus, J, de Zeeuw, P.T., & Lub, J. 1989, *A&A*, 216, 44
- Fitzpatrick, E.L. 1997, *ApJ*, 482, L199
- Gensheimer, P.D., Mauersberger, R., & Wilson, T.L. 1996, *A&A*, 314, 281
- Gredel, R., van Dishoeck, E.F., & Black, J.H. 1991, *A&A*, 251, 625
- Gredel, R., van Dishoeck, E.F., & Black, J.H. 1993, *A&A*, 269, 477
- Jacq, T., Henkel, C., Walmsley, C.M., Jewell, P.R., & Baudry, A. 1988, *A&A*, 199, L5
- Kriss, G.A. 1994 in *ASP Conference Series*, Vol. 61, *Astronomical Data Analysis Software and Systems III*, eds. D.R. Crabtree, R.J. Hanisch, and J. Barnes, p. 437
- Morton, D.C. 1991, *ApJS*, 77, 119
- Neufeld, D.A., & Kaufman, M.J. 1993, *ApJ*, 418, 263
- Neufeld, D.A., Lepp, S., & Melnick, G.J. 1995, *ApJS*, 100, 132
- Riegel, K.W., & Crutcher, R.M. 1972, *A&A*, 18, 55
- Smith, D., & Spañel, P. 1993, *Chem. Phys. Lett.*, 211, 454
- Smith, P.L., Yoshino, K., Griesinger, H.E., Black, J.H. 1981, *ApJ*, 250, 166
- Smith, P.L., & Parkinson, W.H. 1978, *ApJ*, 223, L127

- Snow, T.P., Black, J.H., van Dishoeck, E.F., Burks, G., Crutcher, R.M., Lutz, B.L., Hanson, M.M., & Shuping, R.Y. 1996, ApJ, 465, 245
- Snow, T.P. 1976, ApJ, 204, L127
- Spaans, M. 1996, A&A, 307, 271
- Spaans, M., & van Dishoeck, E.F. 1997, A&A, 323, 953
- van Dishoeck, E.F., Black, J.H., Phillips, T.G., & Gredel, P. 1991, ApJ, 366, 141
- van Dishoeck, E.F., & Black, J.H. 1988, ApJ, 334, 771
- van Dishoeck, E.F., & de Zeeuw, P.T. 1984, MNRAS, 206, 383
- van Dishoeck, E.F., & Helmich, F.P. 1996, A&A, 315, L177
- Vejby-Christensen, L., Andersen, L.H., Heber, O., Kella, D., Pedersen, H.B., Schmidt, H.T., & Zajfman, D. 1997, ApJ, 483, 531
- Williams, T.L., Adams, N.G., Babcock, L.M., Herd, C.R., & Geoghegan, M. 1996, MNRAS, 282, 413
- Zmuidzinas, J., Blake, G.A., Carlstrom, J., Keene, J., Miller, D., Schilke, P., and Ugras, N.G. 1995, in *Proceedings of the Airborne Astronomy Symposium 9 on the Galactic Ecosystem: From Gas to Stars to Dust*, ed. M.R. Haas, J.A. Davidson & E.F. Erickson (San Francisco: ASP), p. 33

Table 1
Velocities, b values, Equivalent Widths,
and Column Densities of Mg II for HD 154368

V_{helio} (km s $^{-1}$)	b (km s $^{-1}$)	EW (mÅ) (λ 1239.9)	EW (mÅ) (λ 1240.4)	log N (cm $^{-2}$)
-36.2	1.2	1.9	1.3	13.7
-26.7	1.3	1.8	1.2	13.6
-16.9	2.7	19.6	13.3	15.0
-1.1	2.9	39.3	26.9	16.2
+10.3	1.5	3.6	2.4	14.0

Figure Captions

- Fig. 1a – GHRS Echelle spectrum of HD 154386, showing the spectral region around the expected position of the 1239.382 Å, 1239.728 Å and 1240.153 Å water features. Vertical lines show the line wavelengths for the heliocentric velocity of the main Mg II component. The labels $1_{01} - 2_{11}$, $0_{00} - 1_{10}$ and $1_{01} - 1_{11}$ denote transitions of the C 1B_1 -X 1A_1 band of water.
- Fig. 1b – GHRS Echelle spectrum of HD 154386, showing the spectral region around the expected position of the 1239.382 Å water feature. The vertical line shows the line wavelength for the heliocentric velocity of the main Mg II component.
- Fig. 2 – GHRS Echelle spectrum of the λ 1239.925, 1240.395 Mg II doublet towards HD 154368. The dashed curves indicate the fits.
- Fig. 3 – GHRS Echelle spectrum of the λ 1237.059 Ge II line towards HD 154368. The vertical line shows the line wavelength for the heliocentric velocity of the main Mg II component.
- Fig. 4a – Depth dependence of some major atomic and molecular species for a model of the translucent cloud in front of HD 154368, with $n_H = 325 \text{ cm}^{-3}$, $I_{UV} = 3$, $A_V = 2.65 \text{ mag}$ and the depletions of Snow et al. (1996).
- Fig. 4b. – Variation of the gas kinetic temperature for the model in Fig 4a.
- Fig. 5 – Model grid for the column densities of OH and H₂O as functions of ζ and $f(\text{H}_2\text{O})$. The solid curves are for $f(\text{O}) = 0$ and the dashed curve is for $f(\text{O}) = 0.3$ (see text). The cross shows the observed OH column density (Black 1997, private communication) and our 3σ upper limit on the H₂O column density.













